

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/305299573>

AN UPDATED REVIEW ON CATHARANTHUS ROSEUS: PHYTOCHEMICAL AND PHARMACOLOGICAL ANALYSIS

Article *in* Indian Research Journal of Pharmacy and Science · July 2016

CITATIONS

0

READS

1,699

5 authors, including:



Asma Nisar

Universiti Malaysia Perlis

5 PUBLICATIONS 1 CITATION

[SEE PROFILE](#)



Awang Soh Mamat

Universiti Malaysia Perlis

50 PUBLICATIONS 38 CITATIONS

[SEE PROFILE](#)



Mohd Irfan Hatim Mohamed Dzahir

Universiti Malaysia Perlis

27 PUBLICATIONS 220 CITATIONS

[SEE PROFILE](#)



Muhammad Syarhabil Ahmad

Universiti Malaysia Perlis

78 PUBLICATIONS 185 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Polyherbal Formulation in Malaysia [View project](#)




Supercritical carbon dioxide [View project](#)

All content following this page was uploaded by [Muhammad Shahzad Aslam](#) on 14 July 2016.

The user has requested enhancement of the downloaded file.

REVIEW



AN UPDATED REVIEW ON *CATHARANTHUS ROSEUS*: PHYTOCHEMICAL AND PHARMACOLOGICAL ANALYSIS

Asma Nisar*, Awang Soh Mamat, Md Irfan Hatim, Muhammad Shahzad Aslam*, Muhammad Syarhabil Ahmad

School of Bioprocess Engineering, Universiti Malaysia Perlis, Kompleks Pusat Pengajian, Jejawi 3, 02600, Arau, Perlis, Malaysia

Submitted on: 28.06.2016;

Revised on: 09.07.2016;

Accepted on: 10.07.2016

ABSTRACT:

Catharanthus roseus commonly known as evergreen herb is one of famous medicinal herb in the field of cancer treatment. Many famous phytochemicals such as vincristine and vinblastine were isolated from this medicinal plant. It has many pharmacological properties such as anti-oxidant, anti-microbial, anti-diabetic, wound healing, anti-ulcer, hypotensive, antidiarrhoeal, hypolipidemic and memory enhancement. Alkaloids are one of major phytochemicals responsible for its anti-cancer properties followed by phenolic compounds such as flavonoids. The purpose of the current study is to document updated data about its traditional uses, isolated bioactive compounds and pharmacological activities reported.

KEYWORDS: *Catharanthus roseus*; Phytochemicals; Pharmacological activities; Medicinal herbs; Traditional medicine.

Corresponding author: Asma Nisar

E-mail address: asmanisar1234@gmail.com

Indian Research Journal of Pharmacy and Science; 9(2016) 631-653


Journal Home Page: <https://www.irjps.in>

INTRODUCTION

Medicinal plants have a long history of usage in traditional medicine. Ethno-botanical information on medicinal plants and their usage by indigenous cultures is useful in the conservation of traditional cultures, biodiversity, community health care and drug development [1]. The authentic knowledge of the usage of medicinal plants passed from one generation to another, after refining and addition [2]. The folk recipes are prepared either from the whole plant or from their different parts, like leaf, stem, bark, root, flower, seed, etc. and also from their secondary product such as gum, resins, and latex [3]. In the human body, medicinal plants interact directly or indirectly with the body chemistry by the chemical constituents. Once the active constituents are absorbed into the blood, these constituents derive the required benefits by circulating and influencing

the blood stream [4]. Plants supply minerals, vitamins, and certain hormone precursors in addition to protein and energy to human body [5]. Trace elements have significant roles in combating a variety of human ailments and disease was observed by the study of elements with respect to indigenous medicinal plants [6].

Catharanthus roseus is an evergreen sub herb plant growing to 1 m tall. The leaves are oval to oblong, 2.5- 9.5 cm. long and 1-3.5 cm. broad glossy green hairless with a pale midrib and a short petiole about 1- 1.8 cm. long and they are arranged in the opposite pairs. The flowers are white to dark pink with a dark red center, with a basal tube about 2.5-3 cm. long and a corolla about 2-5 cm. diameter with 5 petal like lobes. The fruit is a pair of follicles about 2-4 cm. long and 3 mm broad [7].




(A)



(B)



(C)



(D)

Figure -1: (A) Catharanthus Roseus (B) Flower (C) Fruit and Seed (D) Leaves**TRADITIONAL USE OF *CATHARANTHUS ROSEUS***

The plant has historically been used to treat a wide assortment of diseases. It was used as folk remedy for diabetes in Europe for centuries [8]. In India, juice from the leaves was used to treat wasp stings. In Hawaii, the plant was boiled to make a poultice to stop bleeding. In china, it was used as an astringent, diuretic and cough remedy [9]. In central and south America, it was used as a homemade cold remedy to ease lung congestion and inflammation. Throughout the Caribbean, an extract from the flowers was used to make a solution to treat eye irritation and infections. It also had a reputation as amagic plant, Europeans thought it could ward off evil spirits, and

the French referred to it as “violet of the sorcerers” Western researchers finally noticed the plant in 1950’s when they learnt of a tea that Jamaicans were drinking to treat diabetes. They discovered that the plant contains a lot of useful alkaloids (130 in all at last count). Some, such as catharanthine, leurosinesulfate, lochnerine, tetrahydroalstonine, vindoline and vindolinine lower blood sugar level, however, others act as haemostatics (arrest bleeding) and two others, vincristine and vinblastine have anticancerous properties. Periwinkle also contains the alkaloids reserpine and serpentine, which are powerful tranquilizers. Table1. Summarized traditional uses of the *Catharanthus roseus* in different-countries.

Table: 1: Traditional Medicinal Uses of *Catharanthus roseus* in various developed and developing countries.

Countries	Uses	Reference
Australia	Hot water extract of dried leaves is taken orally for menorrhagia, diabetes, and extract of root bark is taken orally as afebrifuge.	[10] & [11]
Brazil	The hot water extract of dried whole plant is taken orally by a human for diabetes mellitus.	[12] & [13]
China	Hot water extract of the aerial parts is taken orally as a menstrual regulator.	[9] & [14]
Cook-Island	Decoction of dried leaves used orally to treat diabetes, hypertension and Cancer.	[15]
Dominica	Hot water extract of leaves is taken orally by pregnant woman to combat primary inertia in childbirth and the boiled leaves are used to treat diabetes.	[16]
England	Hot water extract of dried whole plant is taken orally for the curing of diabetes	[17]

Europe	Decoction of dried leaves is taken orally for diabetes mellitus.	[8]
France	Hot water extract of entire plant is taken as an antigelactagogue	[9]
French	Guinea Hot water extract of the entire plant is taken orally as a cholagogue.	[18]
India	The hot water extract of dried entire plant is taken orally by human for cancer. Hot water extract of dried leaves are used orally to Hodgkin's disease. The root the extract is taken orally for menorrhagia.	[14]&[19]
Jamaica	Hot water extract of dried leaves is taken orally for diabetes.	[20]
Kenya	Hot water extract of dried leaves is taken orally for diabetes.	[20]
Mexico	Infusion of whole plant is taken orally for stomach problem.	[21]
Malaysia	Hot water extract of dried leaves is taken orally for diabetes.	[22]
Mozambique	Hot water extract of leaves is taken orally for diabetes and rheumatism and the root extract is used orally as hypotensive and febrifuge	[23]
North Vietnam	Hot water extract of the aerial parts is taken orally as a menstrual regulator	[9]&[14]
Pakistan	Hot water extract of dried ovules is utilized orally for diabetes.	[24]
Peru	Hot water extract of dried whole plant is used orally by human adults for cancers, heart disease and leishmaniasis.	[25]
Philippines	Hot water extract of the root is taken orally by pregnant women to produce abortion.	[26], [27]&[14]
South Africa	Hot water extract of dried leaves is taken orally for menorrhagia and diabetes.	[10]
South Vietnam	Hot water extract of the entire plant is used orally by human adults as an antigelactagogue.	[9], [14]
Taiwan	Decoction of dried whole plant is used orally by human adults to treat diabetes mellitus, and liver disease,	[27]&[28]
Thailand	Hot water extract of dried entire plant is taken orally for diabetes.	[29]
USA	Hot water extract of leaves is smoked as a euphoriant.	[30]
Venda	Water extract of dried root is taken orally for venereal disease.	[31]
Vietnam	Hot water extract of dried aerial parts is taken orally as a drug in Vietnamese traditional medicine, listed in Vietnamese pharmacopeia (1974 Edition).	[32]
West Indies	Hot water extract of leafy stems is taken orally for diabetes.	[33]

PHARMACOLOGICAL ACTIVITIES:

ANTI OXIDANT PROPERTY:

In the last years, oxidative stress-related diseases/disorders have gained a special attention. Metabolic, neurodegenerative, cardiovascular, mitochondrial diseases and even cancer, are among the most frequent [34] & [35]. Numerous studies have been investigating the underlying triggering factors, in order to understand the mechanisms of action of

free radicals, as well as to discover effective substances towards preventing and even reversing the occurrence of oxidative damages [36] & [37]. Antioxidants, both from natural and synthetic sources, have proved to be highly effective to control the magnitude of free radicals production, to prevent its undesirable effects, as well as to support the organism's antioxidant and detoxifying mechanisms [38], [39] & [40]. Phenolic compounds have shown promising antioxidant properties, with its potential

being directly related to the type of solvent used in the extraction, but also with plant origin, growing conditions, harvesting time, and storage conditions [41] & [42]. The study of the antioxidant potential of phenolic extracts derived from plant species is one of the hot topics among the scientific community; however, in vitro studies are the most common [43], [44] & [45].

Catharanthus roseus contains significant amounts of volatile and phenolic compounds including caffeoylquinic acids and flavonol glycosides which are known to possess antioxidant activity. It has an important role in the body's defense system that acts as antioxidants against reactive oxygen species (ROS), which are harmful by forming such products through normal cell aerobic respiration [46]. Accumulation of free radicals can cause pathological conditions such as ischemia, asthma, arthritis, inflammation, neuro-degeneration, Parkinson's diseases, mongolism, aging process and perhaps dementia [47]. The flower petals, seeds and other parts of *Catharanthus roseus* exhibit antioxidant properties. Thus, phenolic compounds have redox properties that act as reducing agents, hydrogen donors, singlet oxygen quenchers or metal chelators. It has multiple applications in foods, cosmetics, and pharmaceutical industries. Besides antioxidant activity, these compounds exhibit anti-allergic, anti-inflammatory, antimicrobial, anti-thrombotic, cardio protective and vasodilatory effects [48].

The anti oxidant potential of the ethanolic extract of the roots of the two varieties of *Catharanthus roseus* namely rosea (pink flower) and alba (white flower) was obtained by using a different system of the assay such as hydroxyl radical scavenging activity, superoxide radical-scavenging activity, DPPH radical-scavenging activity and nitric oxide radical inhibition method. The result obtained proved that the ethanolic extract of the roots of Periwinkle varieties has exhibited the satisfactory scavenging effect in the entire assay in a concentration dependent manner but *Catharanthus roseus* was found to possess more antioxidant activity than that of C. Alba [49].

ANTI CANCER ACTIVITY:

The anticancer alkaloids vinblastine and vincristine are derived from stem and leaf of *Catharanthus roseus*. These alkaloids have growth inhibition effect

to some human tumors. Vinblastine is used experimentally for the treatment of neoplasms and is recommended for Hodgkin's disease and choriocarcinoma. Vincristine another alkaloid is used for leukemia in children. Different percentage of the methanolic crude extracts of *Catharanthus roseus* was found to show the significant anticancer activity against numerous cell types in vitro condition and especially greatest activity was found against the multidrug resistant tumor types. Vinblastine is sold as Velban and Vincristine as Oncovin [50] & [51].

ANTI DIABETIC ACTIVITY:

The ethanolic extracts of the leaves and flower of *Catharanthus roseus* showed a dose-dependent lowering of blood sugar compare to the standard drug, glibenclamide. The hypoglycemic effect has appeared due to the result of the increase glucose utilization in the liver. The aqueous extract was found to lower the blood glucose to about 20% in diabetic rats when compared to that of the dichloromethane and methanol extracts which lowered the blood glucose level to 49-58%. The hypoglycemic effect has appeared due to the result of the increased glucose utilization in the liver. The hypoglycemic activity of alkaloids isolated from *Catharanthus roseus* have been studied pharmacologically and a remedy derived from the plant has been marketed under the proprietary name Vinculin as a treatment for diabetes [52] & [53].

ANTI MICROBIAL ACTIVITY:

Crude extracts from different parts of the plant were tested for anti-bacterial activity. Extract from leaves showed significantly higher efficacy. The anti-bacterial activity of the leaf extract of the plant was checked against microorganism like *Pseudomonas aeruginosa* NCIM2036, *Salmonella typhimurium* NCIM2501, *Staphylococcus aureus* NCIM5021 and was found that the extracts could be used as the prophylactic agent in the treatment of many of the disease [54].

ANTI-ULCER PROPERTY:

Vincamine and Vindoline alkaloids of the plant showed anti-ulcer property. The plant leaves proved for anti-ulcer activity against experimentally induced gastric damage in rats [55].

HYPOTENSIVE PROPERTY:

Extract of leaves of the plant made a significant change in decreasing the blood pressure. The leaves have been known to contain 150 useful alkaloids among other pharmacologically active compounds. Significant antihyperglycemic and hypotensive activity of the leaf extracts (hydroalcoholic or dichloromethane-methanol) have been reported in laboratory animals [56].

ANTI DIARRHEAL PROPERTY:

The anti-diarrheal activity of the plant's ethanolic leaf extract was tested in the Wistar rats with castor oil as experimental diarrhea inducing agent. *Catharanthus roseus* showed a dose dependent inhibition of the castor oil-induced diarrhea [57].

WOUND HEALING PROPERTY:

Rats treated with 100 mg /kg/day of *Catharanthus roseus* ethanol extract had a high rate of wound contraction significantly decreased epithelization period, and showed significant increase in dry weight and hydroxyproline content of the granulation tissue when compared with the controls. Wound contraction together with increased tensile strength and

hydroxyproline content support the use of *Catharanthus roseus* in the management of wound healing [58].

HYPOLIPIDIMIC EFFECT:

Significant anti atherosclerotic activity was observed in a study as suggested by reduction in the serum levels of total cholesterol, triglycerides, LDL-c, VLDLc and histology of aorta, liver and kidney with the leaf juice of *Catharanthus roseus* (Linn.) G. Donn [59].

MEMORY ENHANCEMENT ACTIVITY:

Vinpocetine has been reported to have a variety of actions that would hypothetically be beneficial in Alzheimer's disease (AD). The only study investigating this agent in a well-defined cohort of AD patients found no benefit. Meta-analysis of older studies of vinpocetine in poorly-defined dementia populations concluded that there is insufficient evidence to support its clinical use at this time. Vinpocetine has been well tolerated at doses up to 60 mg/d in clinical trials of dementia and stroke, and no significant adverse events were observed [60].

CHEMICAL CONSTITUENTS**Table: 2. Chemical constituents of *Catharanthus roseus***

S.No	CHEMICAL CONSTITUENTS	REFERENCE
1.	(-)Tabersonine	[61]
2.	10-Geraniol hydroxylase	[62]
3.	1 O-Hydroxy deacetyl akuammiline	[63]
4.	11-Methoxy tabersonine	[64]
5.	16-Epi vindolinine-N(B)-oxide	[65]
6.	16-Epi-19-(s) vindoline-N-oxide	[66]
7.	16-Epi-trans iso-sitsirikine	[67]
8.	16-Epi-Z-iso-sitsirikine	[67]
9.	16-Hydroxy tabersonine	[68]
10.	16-Methoxy tabersonine	[68]
11.	17-Deacetoxy leurosine	[69]
12.	17 -Deactoxy vincal leukoblastine	[69]
13.	19(s) 16-Epi-vindolinine	[70]
14.	19(s)-Vindolinine	[71]
15.	19-20-Cis 16(R) iso-sitsirikine	[72]
16.	19-20-Trans 16(R) iso-sitsirikine	[72]
17.	19-Acetoxy-11-hydroxy tabersonine	[73]

18.	19-Epi, 3-iso ajmalicine	[74]
19.	19-Epi-ajmalicine	[75]
20.	19-Epi-vindolinine	[73]
21.	19-Hydroxy-11-methoxy tabersonine	[73]
22.	20-Epi-vindolinine	[65]
23.	20-Hydroxy tabersonine	[76]
24.	21-Hydroxy cyclochnerine	[72]
25.	21'-Oxo leurosine	[77]
26.	24-Methylene cholesterol	[78]
27.	2-Hydroxy-6-methoxy benzoic acid	[79]
28.	3'-4'-Anhydro-vincalokoblastine	[80]
29.	3-Epi-ajmalicine	[63]
30.	3-Hydroxy voafrine A	[81]
31.	3-Hydroxy voafrine B	[81]
32.	3-Iso-ajmalicine	[65]
33.	4-Deacetoxy vincalokoblastine	[82]
34.	4-Deacetoxy-3'-hydroxy vincalokoblastine	[83]
35.	S' Phosphodiesterase	[84]
36.	7-Hydroxy-indolenine ajmalicine	[63]
37.	Adenine phosphoribosyltransferase	[85]
38.	Adenoside diphosphate	[86]
39.	Adenoside triphosphate	[86]
40.	Adenosine	[87]
41.	Ajmalicine synthetase	[88]
42.	Ajmalicine	[89] & [90]
43.	Ajmaline	[91]
44.	Akuamicine	[92]
45.	Akuammicine	[93] & [94]
46.	Akuammigine	[76]
47.	Akuammiline	[65]
48.	Akuammine	[95]
49.	Alpha amyrrin acetate	[96]
50.	Alpha-3, 4-anhydrovinblastine synthase	[97]
51.	Alstonine	[98] & [99]
52.	Ammocalline	[93]
53.	Ammorosine	[93]
54.	Amotin	[100]
55.	Anthranilate synthetase	[101]
56.	Antirrhine	[74]
57.	Aparicine	[102]
58.	ATP sulfurylase	[103]
59.	Bannucine	[104]
60.	Beta sitosterol	[105]
61.	Calmodulin	[106]
62.	Campesterol	[78]
63.	Cantharantine	[107]
64.	Carosidine	[108]


65.	Carosine	[95]
66.	Cathalanceine	[109]
67.	Catharanhine	[110]
68.	Catharanthamine	[111]
69.	Catharanthine	[89]
70.	Catharanthus roseus alkaloid (Mp 300+)	[112]
71.	Catharanthus roseus alkaloid (MW 336)	[113]
72.	Catharanthus roseus alkaloid B	[113]
73.	Catharanthus roseus alkaloid C	[113]
74.	Catharanthus roseus alkaloid D	[113]
75.	Catharanthus roseus iridoid glucoside (Mp 194-5)	[114]
76.	Catharicine	[108] & [95]
77.	Catharine	[95]
78.	Catharosine	[115]
79.	Cathasterone	[116]
80.	Cathenamine	[117]
81.	Cathindine	[118] & [93]
82.	Cathovaline:	[119]
83.	Cavincidine	[118] & [93]
84.	Cavincine	[95]
85.	Cholesterol	[78]
86.	Choline	[120]
87.	Cinnamate 4-hydroxylase	[121]
88.	Coronaridine	[122]
89.	Deacetoxy vincalokoblastine	[123]
90.	Deacetoxy vindoline	[124] & [68]
91.	Deacetyl aduammiline	[125]
92.	Deacetyl akuammiline	[63]
93.	Deacetyl vincalokoblastine	[126]
94.	Deacetyl vindoline acetyl transferase	[127]
95.	Deacetyl vindoline	[128] & [68]
96.	Dehydro loganin	[87]
97.	De-n-methyl vincalokoblastine	[129]
98.	Deoxy loganin	[114]
99.	Deoxy vincalokoblastine	[130]
100.	Diacylglycerol pyrophosphate	[131]
101.	Dihydro sitsirikine	[95]
102.	Dihydro vindoline	[132]
103.	Dimethyl tryptamine	[133]
104.	Diol pseudo vincalokoblastine	[134]
105.	Epi-vindolinine	[124]
106.	Extensin	[135] & [136]
107.	Fluorocarpamine	[66]
108.	Fluorocarpamine-N-oxide	[66]
109.	Fructose-2-6-bis-phosphate	[137]
110.	Geissoschizine dehydrogenase	[138]

111.	Geissoschizine	[139]
112.	Geraniol	[140]
113.	Glucose	[120]
114.	Glutamine	[120]
115.	Glycoprotein	[141]
116.	Gomaline	[71]
117.	Hemicellulose	[142]
118.	Hirsutidin	[143] & [144]
119.	Horhammericine	[124]
120.	Indole-3-acetic acid	[145]
121.	Iso-fucoesterol	[78]
122.	Iso-leu rosine	[93]
123.	Iso-pent-2-enyl adenine riboside	[146]
124.	Iso-pent-2-enyl adenine riboside-s'-monophosphate	[146]
125.	Iso-pent-2-enyl adenine	[146]
126.	Isositsirikine	[95]
127.	Isositsirikine	[74]
128.	Isovallesiachotamine	[147]
129.	Isovincoside	[148]
130.	Kaempferol	[79]
131.	L-(+)-bornesitol	[149]
132.	Lanceine	[109]
133.	Leurocolombine	[113]
134.	Leurocristine	[150]
135.	Leurosidine	[150]
136.	Leurosidine-N' -B-oxide	[94]
137.	Leurosine	[151]
138.	Leurosine-N-oxide	[152]
139.	Leurosinone	[153]
140.	Leurosivine	[93] & [95]
141.	Linoleic acid	[154]
142.	Lirioresinol B, D-glucoside	[155]
143.	Lochnerallol	[10]
144.	Lochnericine	[105]
145.	Lochneridine	[130]
146.	Lochnerine	[105]
147.	Lochnerinine	[156]
148.	Lochnerivine	[93] & [118]
149.	Lochnerol	[10]
150.	Lochrovicine	[108]
151.	Lochrovidine	[108]
152.	Lochrovine	[108]
153.	Loganic acid	[157]
154.	Loganin	[87]
155.	Maandrosine	[93]
156.	Malic acid	[158]


157.	Malvidin	[143] & [144]
158.	Mannoside	[149]
159.	Mevalonate kinase	[159]
160.	Minovincinine	[65]
161.	Mitraphylline	[64]
162.	Myricyl alcohol	[160]
163.	N-deformylleurocristine	[161]
164.	N-demethyl vincal leukoblastine	[162]
165.	Neoleurocristine	[108] & [95]
166.	Neoleurosidine	[95]
167.	Nor Harman	[119]
168.	Oleanolic acid	[163]
169.	Oleic acid	[136]
170.	Para-coumaric acid	[79]
171.	Pericalline	[164] & [93]
172.	Pericyclivine	[94]
173.	Perimivine	[118]
174.	Perividine	[93]
175.	Perivine	[105]
176.	Perosine	[118] & [93]
177.	Petunidin	[144] & [143]
178.	Phosphodiesterase	[165]
179.	Phytochelatin A	[166]
180.	Pleiocarpamine	[65]
181.	Pleurosine	[95]
182.	Prolyl hydroxylase	[167]
183.	Protocatechuic acid	[79]
184.	Pseudo-vincal leukoblastine diol	[149]
185.	Putrescine	[168]
186.	Quercetin	[79]
187.	Quinone reductase	[169]
188.	Reserpine	[76] & [170]
189.	Rhazimol	[64]
190.	Ricinoleic acid	[171]
191.	Rosamine	[172]
192.	Roseadine	[173]
193.	Roseoside	[87]
194.	Rosicine	[174]
195.	Rovidine	[108]
196.	Secologanic acid	[175]
197.	Secologanin	[157]
198.	Secologanoside	[175]
199.	Serpentine	[176] & [177]
200.	Sitsirikine	[74]
201.	Stigmasterol	[78]
202.	Strictosidine glucosidase I	[178]
203.	Strictosidine glucosidase II	[178]

204.	Strictosidine lactam	[92]
205.	Strictosidine synthase	[179]
206.	Strictosidine synthetase	[180]
207.	Strictosidine	[181]
208.	Sulfotransferase	[103]
209.	Sweroside	[114]
210.	Syringic acid	[79]
211.	Tabersonine	[182]
212.	Tetrahydro alstonine	[183], [184] & [132]
213.	Tetrahydroserpentine	[156]
214.	Tryptamine	[185]
215.	Tryptophan decarboxylase	[186]
216.	Tryptophan synthetase	[101]
217.	Tryptophan	[101]
218.	Tubotaiwine	[187]
219.	Uridine	[188]
220.	Ursolic acid	[189]
221.	Vallesiachotamine	[73]
222.	Vanillic acid	[79]
223.	Vinamidine	[190]
224.	Vinaphamine	[118]
225.	Vinaspine	[108]
226.	Vincadioline	[191]
227.	Vincaleukoblastine	[192] & [193]
228.	Vincaline I	[194]
229.	Vincaline II	[194]
230.	Vincamicine	[130]
231.	Vincarodine	[94]
232.	Vincathicine	[118]
233.	Vincolidine	[195]
234.	Vincoline	[196]
235.	Vincubine	[197]
236.	Vindolicine	[108] & [95]
237.	Vindolidine	[108]
238.	Vindoline	[198], [110] & [170]
239.	Vindolinine	[199]
240.	Vindolinine-N(B)oxide	[65]
241.	Vindolinine-N-oxide	[199]
242.	Vindorosine	[200]
243.	Vinosidine	[93]
244.	Vinsedicine	[108]
245.	Vinsedine	[108]
246.	Virosine	[105] & [93]
247.	Vivaspine	[118]
248.	Yohimbine	[74]
249.	Zeatin glucosyl	[201]
250.	Zeatin riboside-5'-monophosphate	[146]


251.	Zeatin ribosyl	[201]
252.	Zeatin	[201]
253.	Zeatin-9-riboside	[146]




(A) Kaempferol




(B) Malvidin




(C) Para-coumaric acid




(D) Petunidin




(E) Protocatechuic acid




(F) Quercetin




(G) Rutin



(H) Syringic acid



(I) Vincristine



(J) vinblastin

Figure 2: List of some important bioactive compounds

CONCLUSION:

Catharanthus roseus is one of the important medicinal herb with numerous biological properties. Lot of work is still in process to identify new bioactive compounds, understanding the

methodology of transformation of bioactive compounds in one form to another form, new extraction technique such as green extraction and improving the drying method such as solar drying

REFERENCES:

1. M. Ajaib, Z.U.D. Khan, N. Khan, M. Wahab. Ethnobotanical studies on useful shrubs of District Kotli, Azad Jammu & Kashmir, Pakistan. *Pak J Bot*, 42 (2010), pp. 1407–1415.
2. F. Haq, H. Ahmed, M. Alam. Traditional uses of medicinal plants of Nandiar Khuwarr catchment (District Battagram), Pakistan. *J Med Plants Res*, 5 (2011), pp. 39–48.
3. K.Y. Khan, M.A. Khan, R. Niamat, M. Munir, H. Fazal, P. Mazari, *et al.* Element content analysis of plants of genus *Ficus* using atomic absorption spectrometer. *Afr J Pharm Pharmacol*, 5 (2011), pp. 317–321.
4. A. Kolasani, H. Xu, M. Millikan. Evaluation of mineral content of Chinese medicinal herbs used to improve kidney function with chemometrics. *Food Chem*, 127 (2011), pp. 1465–1471.
5. F. Haq, R. Ullah. Comparative determination of trace elements from *Allium sativum*, *Rheum australe* and *Terminalia*

- chebula* by atomic absorption spectroscopy. Int J Biosci, 1 (2011), pp. 77–82.
6. K. Shirin, S. Imad, S. Shafiq, K. Fatima. Determination of major and trace elements in the indigenous medicinal plant *Withania somnifera* and their possible correlation with therapeutic activity. J Saudi Chem Soc, 14 (2010), pp. 97–100.
 7. Monika Sain and Vandana Sharma. *Catharanthus roseus*- A Review of Potential Therapeutics Properties. Int. J. Pure App. Biosci. 1(6): 139-142 (2013).
 8. Swanston-Flatt SK, Day C, Flatt PR, et al. Glycaemia effects of traditional European plant treatments for diabetes studies in normal and streptozotocin diabetic mice. Diabetes Res. 1989; 10: 69-73.
 9. Farnsworth NR: The pharmacognosy of the periwinkles: Vinca and Catharanthus. Lloydia. 1961; 24 (3): 105-138.
 10. Bhandari PR, Mukerji B. *Lochnera rosea* Linn Reichb. Gauhati Ayurvedic Coll Mag. 1959; 8: 1-4.
 11. Webb LJ: Guide to medicinal and poisonous plants of Queensland. CSIR Bull 232, Melbourne. 1984.
 12. Brandao M, Botelho M, Krettli E. Antimalarial Experimental Chemotherapy using Natural Products. Cienc Cult. 1985; 37(7): 1152-1163.
 13. De Mello JF: Plants in traditional medicine in Brazil. J Ethnopharmacol. 1980; 2(1): 49-55.
 14. Virmani OP, Srivastava GN, Singh P. *Catharanthus roseus* – The tropical periwinkle. Indian Drugs. 1978; 15: 231-252.
 15. Holdsworth DK: Traditional medicinal plants of Rarotonga, Cooks Islands Part I. Int J Crude Drug Res. 1990; 28(3): 209-218.
 16. Hodge WH, Taylor D. The ethanobotany of the island Caribes of Dominica. WEBBIA. 1956; 12: 513-644.
 17. Thompson WAR: Herbs that heal. J. Roy Coll. Gen. Pract. 1976; 26: 365-370.
 18. Luu C: Notes on the traditional pharmacopeia of French Guyana. Plant Med Phytother. 1975; 9: 125- 135.
 19. ANON: Ayurvedic drug to fight cancer. Probe. 1985; 24(2): 234-236.
 20. Morrison EYSA, West M. Indian medicinal plants on blood sugar levels in the dog. West Indian Med J. 1982; 31: 194-197.
 21. Johns T, Kokwaro JO, Kimanani EK. Herbal remedies of the Luo of Siaya District, Kenya. Establishing quantitative criteria for consensus. Econ Bot. 1990; 44(3): 369-381.
 22. H.C. Ong, N. Ahmad, P. Milow. Traditional medicinal plants used by the temuan villagers in Kampung Tering, Negeri Sembilan, Malaysia. Ethnop Med, 5 (2011), pp. 169–173.
 23. Amico A: Medicinal plants of Southern Zambesia. Fitoterapia. 1977; 48: 101-139.
 24. Atta-Ur-Rahman: Some approaches to the study of indigenous medicinal plants. Bull Islamic Med. 1982; 2: 562-568.
 25. Ramirez VR, Mostacero LJ, Garcia AE, et al. Vegetales empleados en medicina tradicional Noreruana. Banco Agrario Del Peru and NACL. Univ Trujillo, Trujillo. Peru, 54pp. 1988.
 26. Zaguirre JC: Guide notes of bed-size preparations of most common local (Philippines) medicinal plants. 1944.
 27. ANON: Description of the Philippines. Part I., Bureau of Public Printing, Manila. 1903.
 28. Hsu FL, Cheng JT. Investigation in rats of the antihyperglycaemic effect of plant extracts used in Taiwan for the treatment of diabetes mellitus. Phytother Res. 1992; 6(2): 108-111.
 29. Yang LL, Yen KY, Kiso Y, et al. Antihepatotoxic Actions of Formosan Plant Drugs. J Ethnopharmacol. 1987; 19 (1): 103-110.
 30. Mueller- Oerlinghausen B, Nagmwathana W, Kanchanapee P. Investigation into Thai medicinal plants said to cure diabetes. J Med Assoc Thailand. 1971; 54: 105-111.
 31. Siegel RK: Herbal intoxication. Psychoactive effects from herbal cigarettes, tea and capsules. J Amer Med Assoc. 1976; 236 (5): 473-476.
 32. Arnold HJ, Gulumian M. Pharmacopea of traditional medicine in Venda. J Ethnopharmacol. 1984; 12 (1): 35-74.

33. Nguywen, Van D. List of simple drugs and medicinal plants of value in Vietnam. Proc Seminar of the use of Medicinal Plants in Healthcare. Tokyo 13-17, September, WHO Regional Officer Manila.1977; 65- 83.
34. Halliwell, Barry. "Free radicals and antioxidants: updating a personal view." *Nutrition reviews* 70.5 (2012): 257-265.
35. Singh, Ravindra Pratap, Shashwat Sharad, and Suman Kapur. "Free radicals and oxidative stress in neurodegenerative diseases: relevance of dietary antioxidants." *J Indian Acad Clin Med* 5.3 (2004): 218-225.
36. Espín, Juan Carlos, María Teresa García-Conesa, and Francisco A. Tomás-Barberán. "Nutraceuticals: facts and fiction." *Phytochemistry* 68.22 (2007): 2986-3008.
37. Fernandez-Panchon, M. S., et al. "Antioxidant activity of phenolic compounds: from in vitro results to in vivo evidence." *Critical reviews in food science and nutrition* 48.7 (2008): 649-671.
38. Holst, Birgit, and Gary Williamson. "Nutrients and phytochemicals: from bioavailability to bioefficacy beyond antioxidants." *Current opinion in biotechnology* 19.2 (2008): 73-82.
39. Kapravelou, Garyfallia, et al. "Improvement of the antioxidant and hypolipidaemic effects of cowpea flours (*Vigna unguiculata*) by fermentation: results of in vitro and in vivo experiments." *Journal of the Science of Food and Agriculture* 95.6 (2015): 1207-1216.
40. Valko, Marian, et al. "Free radicals and antioxidants in normal physiological functions and human disease." *The international journal of biochemistry & cell biology* 39.1 (2007): 44-84.
41. Avello, Marcia A., et al. "Variation in phenolic compounds of *Ugni molinae* populations and their potential use as antioxidant supplement." *Revista Brasileira de Farmacognosia* 23.1 (2013): 44-50.
42. Taârit, Mouna Ben, et al. "Physiological changes, phenolic content and antioxidant activity of *Salvia officinalis* L. grown under saline conditions." *Journal of the Science of Food and Agriculture* 92.8 (2012): 1614-1619.
43. Dai, Jin, and Russell J. Mumper. "Plant phenolics: extraction, analysis and their antioxidant and anticancer properties." *Molecules* 15.10 (2010): 7313-7352
44. Larrosa, Mar, et al. "Ellagitannins, ellagic acid and vascular health." *Molecular aspects of medicine* 31.6 (2010): 513-539.
45. Rubió, Laura, María-José Motilva, and María-Paz Romero. "Recent advances in biologically active compounds in herbs and spices: a review of the most effective antioxidant and anti-inflammatory active principles." *Critical reviews in food science and nutrition* 53.9 (2013): 943-953.
46. Salah, Nida, et al. "Polyphenolic flavanols as scavengers of aqueous phase radicals and as chain-breaking antioxidants." *Archives of biochemistry and biophysics* 322.2 (1995): 339-346.
47. Kaur, Sarabjot, and Poonam Mondal. "Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants." *Journal of microbiology and Experimentation* 1.1 (2014): 00005.
48. Garg, Vishal, et al. "Granulocyte colony-stimulating factor mobilizes CD34+ cells and improves survival of patients with acute-on-chronic liver failure." *Gastroenterology* 142.3 (2012): 505-512.
49. Alba Bhutkar MA, Bhise SB. Comparative Studies on Antioxidant Properties of *Catharanthus Rosea* and *Catharanthus*. *International Journal of Pharmaceutical Techniques*, 2011, 3(3): 1551-1556.
50. Banskota AH. Antiproliferative activity of Vietnamese medicinal plants. *Biological Pharmaceutical Bulletin*, 2002, 25(6):753-60.
51. Wang S, Zheng Z, Weng Y. Angiogenesis and antiangiogenesis activity of Chinese medicinal herbal extracts. *Life Science*, 2004, 74(20): 2467-78.
52. Chattopadhyay RR, Sarkar SK, Ganguli S. Hypoglycemic and antihyperglycemic effect

- of leaves of *Vinca rosea* Linn. Indian Journal of Physiology and Pharmacology, 1991, 35: 145-51.
53. Chattopadhyay RR.A comparative evaluation of some blood sugar lowering agents of plant origin. Journal of Ethnopharmacology, 1994, 67:367-72.
54. Prajakta J. Patil, Jai S. Ghosh. Antimicrobial Activity of *Catharanthus roseus* – A Detailed Study. British Journal of Pharmacology and Toxicology, 2010, 1(1): 40-44.
55. Babulova A, Machova J, Nosalova V. Protective action of vinpocetine against experimentally induced gastric damage in rats. Arzneimittel forschung, 2003, 43:981-985.
56. P. P. Pillay, C. P. M. Nair, and T. N. Santi Kumari. *Lochnera rosea* as a potential source of hypotensive and other remedies. Bulletin of Research Institute of the University of Kerala, 1959, 1:51-54.
57. Mithun Singh Rajput, Veena Nair, Akansha Chauhan. Evaluation of Antidiarrheal Activity of Aerial Parts of *Vinca major* in Experimental Animals. Middle-East Journal of Scientific Research. 2011, 7 (5): 784-788.
58. Nayak BS, Anderson M and Pereira LMP. Evaluation of wound-healing potential of *Catharanthus roseus* leaf extract in rats. Fitoterapia, 2007, 78:540-544.
59. Yogesh Patel et al. Evaluation of hypolipidemic activity of leaf juice of *Catharanthus roseus* (Linn.). Acta Poloniae Pharmaceutica - Drug Research, 2011, 68 (6) :927-935.
60. P. Sekar. Vedic clues to memory enhancer. The Hindu, March 21, 1996.
61. Petiard, V., F. Gueritte, N. Langolis and P. Potier. Presence of (-) tabersonine in a tissue culture strain of *Catharanthus roseus* G. Don. Physiol Veg 1980; 18: 711-720.
62. Schiel, O. and J. Berlin. Geraniol-10-hydroxylase of *Catharanthus roseus* and its correlation with indole alkaloid biosynthesis. Planta Med 1986; 1986(1): 422A.
63. Gueritte, F., N. Langlois and V. Petiard. Secondary metabolites isolated from a tissue culture of *Catharanthus roseus*. J Nat Prod 1983; 46(1): 144-148.
64. Atta-ur-Rahman, I. Ali and M. Bashir. Isolation of rhazimol from the leaves of *Catharanthus roseus*. J Nat Prod 1984; 47(2): 389-.
65. Kohl, W., B. Witte and G. Hofle. Alkaloids from *Catharanthus roseus* tissue cultures. II. Z Naturforsch Ser B 1981; 36: 1153-1162.
66. Rahman, A. U. and M. Bashir. Isolation of new alkaloids from *Catharanthus roseus*. Planta Med 1983; 49{2}: 124-125.
67. Mukhopadhyay, S., A. El-Sayed, G. A. Handy and G. A. Cordell. Catharanthus alkaloids. XXXVII. 16-Epi-z-isositsirikine, a monomeric indole alkaloid with antineoplastic activity from *Catharanthus roseus* and *Rhazya stricta*. J Nat Prod 1983; 46(3): 409-413.
68. De Luca, V., N. Brisson, J. Balsevich and W. G. W. Kurz. Regulation of vindoline biosynthesis in *Catharanthus roseus*: Molecular cloning of the first and last steps in biosynthesis. Primary and Secondary Metabolism of Plant Cell Cultures II. 1989; 154-161.
69. De Bruyn, A., L. De Taeye, R. Simonds, M. Verzele and C. De Pauw. Alkaloids from *Catharanthus roseus*. Isolation and identification of 17 desacetoxylvindoline and 17 -desacetoxyleurosine. Bull Soc Chim Belg 1982; 91(1): 75-85.
70. Atta-Ur-Rahman, M. Bashir, S. Kaleem and T. Fatima. 16-Epi-19-svindolinine, an indole alkaloid from *Catharanthus roseus*. Phytochemistry 1983; 22(4): 1021-1023.
71. Atta-ur-Rahman and I. Ali. 13-C NMR Spectroscopic studies on gomaline and rosamine. Fitoterapia 1986; 57(6): 438-440.
72. Kohl, W., B. Witte, W. S. Sheldrick and G. Hofle. Indole alkaloids from *Catharanthus roseus* tissue cultures. IV. 16r-19 ,20-e-isositsirikin, 16e-19 ,20-z- isositsirikin and 21-hydroxycyclolochnerin. Planta Med 1984; 1984{3}: 242-244.

73. Kurz, W. G. W., K. B. Chatson, F. Constabel, J. P. Kutney, L. S. L. Choi, P. Kolodziejczyk, S. K. Sleight and K. L. Stuart. The production of catharanthine and other indole alkaloids by cell suspension cultures of *Catharanthus roseus*. *Planta Med* 1980; 39: 284A.
74. Kohl, W., B. Witte and G. Hofle. Alkaloids from *Catharanthus roseus* tissue cultures. 3. *Z Naturforsch Ser B* 1982; 37: 1346-1351.
75. Treimer, J. F. and M. H. Zenk. Enzymic synthesis of corynanthe-type alkaloids in cell cultures of *Catharanthus roseus*: Quantation by radioimmunoassay. *Phytochemistry* 1978; 17: 227-231.
76. Kohl, W., B. Witte and G. Hofle. Quantitative and qualitative HPLC analysis of indole alkaloids from *Catharanthus roseus* cell cultures. *Planta Med* 1983; 47(3): 177-182.
77. El-Sayed, A. G. A. Handy and G. A. Cordell. *Catharanthus* alkaloids. XXXIII. 21'-Oxoleurosine from *Catharanthus roseus* (Apocynaceae). *J Nat Prod* 1980; 43: 157-161.
78. Duperon, P., J. P. Allais and A. Dupaix. Comparison of the sterol content of tonoplast and microsomal fractions from *Catharanthus roseus* suspensioncultured cells. *Plant Physiol Biochem (Paris)* 1992; 30(4): 495-498.
79. Daniel, M. and S. D. Sabnis. Chemotaxonomical studies on Apocynaceae. *Indian J Exp Bioi* 1978; 16(4): 512-513.
80. ANON. Isolating alkaloid compounds from *Vinca rosea*. Patent-Belg-867,670 1978.
81. Fahn, W., V. Kaiser, H. Schubel, J. Stockigt and B. Daniel. *Catharanthus roseus* enzyme mediated synthesis of 3- hydroxyvoafrine A and B - A simple route to the voafrines. *Phytochemistry* 1990; 29(1): 127-133.
82. Neuss, N. and A. J. Barnes. 4-Deacetoxyvinblastine. Patent-US-3,954, 773 1976.
83. Tafur, S. S. 4-Desacetoxy-3-hydroxyvinblastine. Patent-US-3,944,554 1976.
84. Ukita, M., A. Furuya, H. Tanaka and M. Misawa. S'Phosphodiesterase formation by cultured plant cells. *Agr Bioi Chern* 1973; 37: 2849-2854.
85. Hirose, F. and H. Ashihara. Adenine phosphoribosyltransferase activity in mitochondria of *Catharanthus rose us* cells. *Z Naturforsch Ser C* 1982; 37(11/12): 1288-1289.
86. Shimazaki, A., F. Hirose and H. Ashihara. Changes in adenine nucleotide levels and adenine salvage during the growth of *Vinca rosea* cells suspension culture. *Z Pflanzenphysiol* 1982; 106: 191-198.
87. Bhakuni, D. S., P. P. Joshi, H. Uprety and R. S. Kapil. Roseoside - A C-13 glycoside from *Vinca rosea*. *Phytochemistry* 1974; 13: 2541-2543.
88. Scott, A. I., S. L. Lee and W. Wan. Indole alkaloid biosynthesis: Partial purification of "ajmalicine synthetase" from *Catharanthus roseus* tissue cultures. *Heterocycles* 1977; 6: 1552.
89. Hars, Y. and F. Ito. Manufacture of indole alkaloids with plant tissue or organ culture. Patent-Japan Kokai Tokkyo Koho-04 1992; 91-785: 7pp.
90. Singh, J., K. L. Handa, P. R. Rao and C. K. Atal. Recovery of ajmalicine (raubasine) from *Vinca rosea*. *Res Ind* 1978; 23: 166-167.
91. Kamata, H. Production of vinca-alkaloid. Patent-Japan Kokai Tokkyo Koho-61 1986; 274,694: 6pp.
92. Constabel, F., S. Rambold, K. B. Chatson, W. G. M. Kurz and J. P. Kutney. Alkaloid production in *Catharanthus rose us* (L.) G. Don. VI. Variation in alkaloid spectra of cell lines derived from one single leaf. *Plant Cell Rep* 1981; 1(1): 3-5.
93. Svoboda, G. H., A. T. Oliver and D. R. Bedwell. Alkaloids of *Vinca rosea* (*Catharanthus roseus*). XIX. Extraction and characterization of root alkaloids. *Lloydia* 1963; 26: 141.

94. Mukhopadhyay, S. and G. A. Cordell. Catharanthus alkaloids. XXXVI. Isolation of vincalukoblastine (VLB) and periformylone from *Catharanthus trichophyllus* and pericyclivine from *Catharanthus roseus*. *J Nat Prod* 1981; 44: 335-339.
95. Willaman, J.J. and H. L. Li. Alkaloidbearing plants and their contained alkaloids, 1957-1968. *Lloydia* 1970; 33S: 1-286.
96. Khan, N. and D. P. Chakraborty. Alpha amyrrin acetate from *Catharanthus roseus* Linn. *J Indian Chem Soc* 1981; 58: 628-629.
97. Sottomayor, M., M. Lopez Serrano, F. DiCosmo and A. RosBarcelo. Purification and characterization of alpha- 3' ,4'-anhydrovinblastine synthase (peroxidase-like) from *Catharanthus roseus* (L.) G. Don. *FERBS lett* 1998; 428(3): 299-303.
98. Pillay, P. P. and T. N. Santha Kumari. The occurrence of alstonine in *Lochnera rosea*. *J Sci Ind Res-B* 1961; 20: 458.
99. Shimuzu, M. and F. Uchamaru. Isolation of alkaloids from *Vinca* (*Lochnera*) *rosea*. *Chem Pharm Bull* 1958; 6: 324.
100. Sedakova, L. A, I. K. Bukharova, I. D. Treshchalina and A B. Syrkin. Antitumor and toxic effects of arnotin. *Eksp Onkol* 1987; 9(5): 76-77.
101. Scott, A. I., H. Mizukami and S. L. Lee. Characterization of a 5-methyltryptophan resistant strain of *Catharanthus roseus* cultured cells. *Phytochemistry* 1979; 18: 795-798.
102. Rojas, M. C. N. and M. C. A. Cuellar. Comparative microbiological studies of the alkaloids of *Catharanthus roseus* and other related compounds. *Rev Cubana Farm* 1981; 15(2): 131-138.
103. Schwenn, J. D., H. El-Shagi, A. Kemena and E. Petrak. On the role of sulfotransferases in assimilatory sulfate reduction by plant suspension cultures. *Planta* 1979; 144: 419-425.
104. Atta-ur-Rahman, I. Ali and M. I. Chaudhary. Bannucine-A new dihydroindole alkaloid from *Catharanthus roseus* (L) G. Don. *J Chem Soc Perkin Trans I* 1986(6); 923-926.
105. Svoboda, G. H., N. Neuss and M. Gorman. Alkaloids of *Vinca rosea* Linn. (*Catharanthus roseus* G. Don). V. Preparation and characterization of alkaloids. *J Amer Pharm Ass Sci Ed* 1959; 48: 659-666.
106. Radvany, L. G. and F. Di Cosmo. Purification and characterization of calmodulin from cultured cells of *Catharanthus roseus*. *Phytochem Anal* 1991; 2(6): 241-252.
107. Verzele, M., L. De Taeye, J. Van Dyck, G. De Decker and C. De Pauw. High-performance liquid chromatography of *Vinca rosea* alkaloids and the correlation of plate height and molecular weight. *J Chromatogr* 1981; 214: 95-99.
108. Svoboda, G. A. The alkaloids of *Catharanthus roseus* G. Donn (*Vinca rosea* L.). *Current Topics in Plant Science*. J. E. Gunckel (Ed.), Academic Press, New York 1969; 303-335.
109. Patterson, B. D. and D. P. Carew. Growth and alkaloid formation in *Catharanthus roseus* tissue cultures. *Lloydia* 1969; 32: 131.
110. Leete, E., A. Ahmad and I. Kompis. Biosynthesis of the *Vinca* alkaloids. I. Feeding experiments with tryptophan- 2-C¹⁴ and acetate-1-C¹⁴. *J Amer Chem Soc* 1965; 87: 4168.
111. El-Sayed, A. and G. A. Cordell. Catharanthus alkaloids. XXXIV. Catharanthamine, a new antitumor bisindole alkaloid from *Catharanthus roseus*. *J Nat Prod* 1981; 44: 289-293.
112. Weiss, S. G. Antitumor principles of *Unum album* and *Catharanthus roseus*. *Diss Abstr Int* 1974; 35: 2669.
113. Madati, P. J., H. A. Pazi and A. Ernest. Phytochemical investigation of *Catharanthus roseus* growing wild in East Africa. *J Afr Med PI* 1979; 2: 1.
114. Bhakuni, O. S. and R. S. Kapil. Monoterpene glycosides from *Vinca rosea*. *Indian J Chem* 1972; 10: 454.

115. Moza, B. K. and J. Trojanek. Catharosine, a new alkaloid from *Catharanthus roseus*. *Chern Ind (London)* 1965; 1965: 1260.
116. Fujioka, S., T. Inoue, S. Takatsuto, T. Yanagisawa, T. Yokota and A. Sakurai. Identification of a new brassinosteroid, cathasterone, in cultured cells of *Catharanthus roseus* as a biosynthetic precursor of teasterone. *Biosci Biotech Biochem* 1995; 59(8): 1543-1547.
117. Stockigt, J., H. P. Husson, C. Kan-Fan and M. H. Zenk. Cathenamine, a central intermediate in the cell free biosynthesis of ajmalicine and related indole alkaloids. *Chern Commun* 1977; 164-166.
118. Svoboda, G. H. and A. J. Barnes, Jr. Alkaloids of *Vinca rosea* Linn. (*Catharanthus roseus* G. Don) XXXIV. Vinaspine, vincathicine, rovidine, desacetyl vlb and vinphamine. *J Pharm Sci* 1964; 53: 1227.
119. Atta-ur-Rahman, S. Hasan and M. R. Qulbi. Beta-carboline from *Catharanthus roseus*. *Planta Med* 1985; 1985(3): 287.
120. Schripsema, J. and R. Verpoorte. Regulation of indole alkaloid biosynthesis in *Catharanthus roseus* cell suspension cultures, investigated with ¹H-NMR. *Planta Med Suppl* 1992; 58(1): A608.
121. Hotze, M., G. Schroder and J. Schroder. Cinnamate 4-hydroxylase from *Catharanthus roseus*, and a strategy for the functional expression of plant cytochrome P450 proteins as translational fusions with P450 reductase in *Escherichia coli*. *FEBS Lett* 1995; 374(3): 345-350.
122. De Taeye, L., A. De Bruyn, C. De Pauw and M. Verzele. Alkaloids of *Vinca rosea*: Isolation and identification of coronaridine. *Bull Soc Chim Belg* 1981; 90(1): 83-87.
123. Neuss, N., A. J. Barnes and L. L. Huckstep. *Vinca* alkaloids. XXXV. Desacetoxyvinblastine: A new minor alkaloid from *Vinca rosea* (*Catharanthus roseus*). *Experientia* 1975; 31: 18.
124. Balsevich, J., L. R. Hogge, A. J. Berry, D. E. Games and I. C. Mylchreest. Analysis of indole alkaloids from leaves of *Catharanthus roseus* by means of supercritical fluid chromatography/ mass spectrometry. *J Nat Prod* 1988; 51 (6): 1173-1177.
125. Rahman, A-U and S. Malik. Isolation of isoallesiachotamine from legumes of *Rhazya stricta*. *J Nat Prod* 1984; 47(2): 388-389.
126. Nagy-Turak, A. and Z. Vegh. Extraction and in situ densitometric determination of alkaloids from *Catharanthus roseus* by means of overpressured layer chromatography on amino-bonded silica layers. 1. Optimization and validation of the separation system. *J Chromatogr A* 1994; 668(2): 501-507.
127. De Luca, V., J. Balsevich and M. G. W. Kurz. Purification and properties of deacetylindoline acetyltransferase from *Catharanthus roseus*. *Abstr Internat Res Cong Nat Prod Coll Pharm Univ, Chapel Hill NC July 7-12, 1985; Abstr-103*.
128. Groger, D., K. Stolle and C. P. Falshaw. Isolation of desacetylindoline from *Catharanthus roseus*. *Naturwissenschaften* 1965; 52: 132.
129. Jovanovic, K., K. Szasz, G. Fekete, E. Bittner, E. Dezseri and J. Eles. Isolation of vincristine and n-demethylvinblastine from *Vinca rosea*. *Patent-Ger Offen-2,259,388* 1974.
130. Svoboda, G. H., M. Gorman, N. Neuss and A. J. Barnes, Jr. Alkaloids of *Vinca rosea* (*Catharanthus roseus*). VIII. Preparation and characterization of new minor alkaloids. *J Pharm Sci* 1961; 50: 409.
131. Wissing, J. B. and H. Behrbohm. Diacylglycerol pyrophosphate, a novel phospholipid compound. *Febs Lett* 1993; 315(1): 95-99.
132. Roger, D. and K. Stolle. A new alkaloid from *Catharanthus roseus*. *Naturwissenschaften* 1964; 51: 637
133. Kutney, J. P. Studies in plant tissue culture - potentially important sources of clinically important anti-cancer agents. *Proc F E C S Third Int Conference on Chemistry and Biotechnology of Biologically Active Natural Products, Vol 1 PPI68-179, VCH Publisher Inc NY, NY 1973; 1: 168-179*.

134. Tafur, S. S., W. E. Jones, D. E. Dorman, E. E. Logsdon and G. H. Svoboda. Alkaloids of *Vinca rosea* (*Catharanthus roseus*). XXXVI. Isolation and characterization of new dimeric alkaloids. *J Pharm Sci* 1975; 64: 1953.
135. Tanaka, M., H. Shibata, K. Sato, S. Tanada and T. Uchida. Biosynthesis of extensin in suspension-cultured cells of *Vinca rosea*. *Plant Tissue Cult Proc Int Congr Plant Tissue Cell Cult 5th* 1982; 1982: 39--40.
136. Dolya, V. S., N. V. Svinidze, Y. I. Kornievskii and T. R. Gokitidze. Fatty oils of four plants of the Subtropics. *Farm Zh (Kiev)* 1985(2); 76-n.
137. Ashihara, H. changes in fructose-2,6-biphosphate level during the growth of suspension cultured cells of *Catharanthus roseus*. *Z Naturforsch Ser C* 1986; 41(5/6): 529-531.
138. Rueffer, M., C. Kan-Fan, H. P. Husson, J. Stockigt and M. H. Zenk. 4,21-Dehydrogeissoschizine, an intermediate in heteroyohimbine alkaloid biosynthesis. *Chern Commun* 1979; 1016-1018.
139. Scott, A. I and S. L. Lee. Biosynthesis of the indole alkaloids. A cell-free system from *Catharanthus roseus*. *J Amer Chem Soc* 1975; 97: 6906.
140. Deus, B. Cell cultures of *Catharanthus roseus*. Production of Natural Compounds by Cell Culture Methods. *Proc Int Symp Plant Cell Cult, Ges Strahlen Umweltforsch, Munich (AI Fermann, A W, Reinhard, EC, EDS)* 1978; 118.
141. Tanaka, M. and T. Uchida. Heterogeneity of hydroxypyroline-containing glycoproteins in protoplasts from a *Vinca rosea* suspension culture. *Plant Cell Physiol* 1979; 20: 1295-1306.
142. Takeuchi, Y. and A. Komamine. Turnover of cell wall polysaccharides of a *Vinca rosea* suspension culture. I. Synthesis and degradation of cell wall components. *Physiol Plant* 1980; 48: 271-277.
143. Carew, D. P. and R. J. Krueger. Anthocyanidins of *Catharanthus roseus* callus cultures. *Phytochemistry* 1976; 15: 442A.
144. Milo, J. Flower color inheritance and shoot and ajmalicine yield components in successive developmental stages of pure lines and F-1 hybrids in *Catharanthus roseus* (L.) G. Don. Thesis-MS-Hebrew University 1981; 1981: 74 pp.
145. Bandyopadhyay, M. N., B. B. Mukherjee and S. N. Ganguly. IAA from the pollen grains and style stigma segments of *Catharanthus roseus* (Linn.) G. Don. *Physiol Sex Reprod Flowering Plants Int Symp 1st* 1976 1978; 102-104.
146. Scott, I. M., B. A. McGaw, R. Horgan and P. E. Williams. Biochemical studies on cytokinins in *Vinca rosea* crown gall tissue. *Plant Growth Subst Proc Int Conf 11th* 1982; 1982: 165-174.
147. Petiard, Y., D. Courtois, F. Gueritte, N. Langlois and B. Mompon. New alkaloids in plant tissue cultures. *Proc 5th Inti Cong Plant Tissue & Cell Culture* 1982; 309-310.
148. Stoeckigt, J. and M. H. Zenk. Strictosidine {isovincoside}: The key intermediate in the biosynthesis of monoterpene indole alkaloids. *Chern Commun* 1977; 1977(18): 646-648.
149. Rizk, A. F. M. The Phytochemistry of the Flora of Qatar, Scientific and Applied Research Centre, University of Qatar, Kingsprint, Richmond, UK 1986; 582pp.
150. Svoboda, G. H., A. J. Barnes Jr. and R. J. Armstrong. Leurosidine and leurocristine and their production. Patent US-3,205,220 1965.
151. Svoboda, G. H. A note on several new alkaloids from *Vinca rosea*. I. Leurosine, virosine, peri vine. *J Amer Pharm Ass Sci Ed* 1958; 47: 834.
152. EI-Sayed, A, G. H. Handy and G. A. Cordell. Further dimeric indole alkaloids from *Catharanthus roseus* (L.) G. Don (Apocynaceae). *J Nat Prod* 1979; 42: 687D.
153. Atta-ur-Rahman, M. Alam, I. Ali, Habib-ur-Rehman and I. Hag. Leurosine: A new binary indole alkaloid from *Catharanthus roseus*. *J Chern Soc Perkin Trans* 11988(8); 2175-2178.

154. Maccarthy, J. J. and P. K. Stumpf. Fatty acid composition and biosynthesis on cell suspension cultures of Glycine max, Catharanthus roseus and Nicotiana tabacum. *Planta (Berl)* 1980; 147: 384-388.
155. Arfmann, H. A., W. Kohl and V. Wray. Effect of 5-azacytidine on the formation of secondary metabolites in Catharanthus roseus cell suspension cultures. *Z Naturforsch Ser C* 1985; 40: 21-25.
156. Moza, B. K. and J. Trojanek. On alkaloids. VII. New alkaloids from Catharanthus roseus. *Collect Czech Chern Cornrnun* 1963; 28: 1419.
157. Guarnaccia, R., L. Botta and C. J. Coscia. Biosynthesis of acidic iridoid monoterpene glucosides in *Vinca rosea*. *J Amer Chem Soc* 1974; 96: 7079.
158. Marigo, G., H. Bouyssou and B. M. Mal. Vacuolar efflux of malate and its influence on nitrate accumulation in Catharanthus roseus cells. *Plant Sci* 1985; 39(2): 97-103.
159. Schulte, A E., R. van der Heijden and R. Verpoorte. Purification and characterization of mevalonate kinase from suspension-cultured cells of Catharanthus roseus (L.) G. Don. *Arch Biochem Biophys* 2000; 378(2): 287-98.
160. Tangkongchitr, U. Extraction of organic compounds from *Vinca rosea* Linn. Master Thesis 1973; 43pp.
161. Simonds, R., A. De Bruyn, L. De Taeye, M. Verzele and C. De Pauw. Ndeformyl vincristine: A bisalkaloid from Catharanthus roseus. *Planta Med* 1984; 1984{3}: 274-276.
162. Jovanovics, K., K. Szasz, G. Pekete, E. Bittner E. Dezseri and J. Eles. Increasing the yield of vincristine when separating vincristine from *Vinca rosea* drug. *Patent-S Afr-* 08,534 1973; 72.
163. Ali, M. E., N. Huq, A. K. M. M. Rahman, A. Moitra and A. Rahman. Chemical investigations on *Vinca rosea* Linn. (Nayantara). *Bangladesh J Sci Ind Res* 1979; 14: 354-358.
164. Rojas, N. M. and A. Cuellar. Catharanthus roseus G. Don. I. Microbiological study of its alkaloids. *First Latin American & Caribbean Symposium on Pharmacologically Active Natural Products, Cuba 1982 UNESCO 1982*; 194.
165. Furuya, A., M. Ukita, Y. Kotani, M. Misawa and H. Tanaka. RNA hydrolysis to 5'-ribonucleotides by phosphodiesterase from plant callus. *Patent-Japan Kokai-73* 33,903 1973.
166. Grill, E., E. L. Winnacker and M. H. Zenk. Phytochelatins: The principal heavy-metal complexing peptides of higher plants. *Science* 1985; 230(4726): 674-676.
167. Tanaka, M., H. Shibata and T. Uchida. A new prolyl hydroxylase acting on poly-L-proline, from suspension cultured cells of *Vinca rosea*. *Biochim Biophys Acta* 1980; 616: 188-198.
168. Zhou, X. H., R. Minocha and S. C. Minocha. Physiological response of suspension cultures of Catharanthus roseus to aluminum: Changes in polyamines and inorganic ions. *J Plant Physiol* 1995; 145(3): 277-284.
169. Spitsberg, V. L. and C. J. Coscia. Quinone reductases of higher plants. *Eur J Biochem* 1982; 127(1): 67-70.
170. Erdelsky, K. and L. Holickova. Cultivation of plant tissue culture of *Vinca rosea* L. and its alkaloid contents. *Acta Fac Rerum Nat Univ Comeniana Physiol Plant* 1978(15); 1-10.
171. Garg, S. P., M. R. K. Sherwani, A. Arora, R. Agarwal and M. Ahmad. Ricinoleic acid in *Vinca rosea* seed oil. *J Oil Technol Ass India (Bombay)* 1987; 19(3): 63-64.
172. Atta-Ur-Rahman, I. Ali, M. Bashir and M. I. Choudhary. Isolation and structure of rosamine - A new pseudoindoxyl alkaloid from Catharanthus roseus. *Z Naturforsch Ser B* 1984; 39{9}: 1292-1293.
173. El-Sayed, A, G. A Handy and G. A Cordell. Catharanthus alkaloids. XXXVIII. Confirming structural evidence and antineoplastic activity of the bisindole alkaloids leurosine-n'-oxide {pleurosine}, roseadine and vindolicine from Catharanthus roseus. *J Nat Prod* 1983; 46{4}: 517-527.
174. Atta-Ur-Rahman, J. Fatima and K. Albert. Isolation and structure of roscicine from

- Catharanthus roseus. *Tetrahedron Lett* 1984; 25{52}: 6051-6054.
175. Guarnaccia, R. and C. J. Coscia. Occurrence and biosynthesis of secologanic acid in *Vinca rosea*. *J Amer Chem Soc* 1971; 93: 6320.
176. Kutney, J. P. Studies on plant tissue cultures. *Heterocycles* 1981; 15: 1405-1431.
177. Wahl, J. Airlift fermenter for growing plant cell cultures. *Git Fachz Lab* 1979; 23: 169-171.
178. Hemscheidt, T. and M. H. Zenk. Glucosidases involved in indole alkaloid biosynthesis of *Catharanthus* cell cultures. *Febs Lett* 1980; 110: 187-191.
179. Treimer, J. F. and M. H. Zenk. Purification and properties of strictosidine synthase, the key enzyme in indole alkaloid formation. *Eur J Biochem* 1979; 101: 225-233.
180. Mizukami, H., H. Norlov, S. L. Lee and A. I. Scott. Purification and properties of strictosidine synthetase (an enzyme condensing tryptamine and secologanin) from *Catharanthus roseus* cultured cells. *Biochemistry* 1979; 18: 3760-3763.
181. Chen, T. H. H., K. K. Kartha, N. L. Leung, W. G. W. Kurz, K. B. Chatson and F. Constabel. Cryopreservation of alkaloid-producing cell cultures of periwinkle (*Catharanthus roseus*). *Plant Physiol* 1985; 75: 726-731.
182. Auriola, S., T. Naaranlahti and S. P. Lapinjoki. Determination of *Catharanthus roseus* alkaloids by high-performance liquid chromatography isotope dilution thermospray-mass spectrometry. *J Chromatogr* 1991; 554(1/2): 227-231.
183. Shimizu, M. and F. Uchamaru. The isolation of alkaloids from *Vinca (Lochnera) rosea*. *Chem Pharm Bull* 1959; 7: 713.
184. Ali, I. The alkaloids in flowers of *Catharanthus roseus* (L) G. Don. *Gomal Univ J Res* 1990; 100): 27-31.
185. Islas, I., V. M. Loyola-Vargas and M. De Lourdes Miranda-Ham. Tryptophan decarboxylase activity in transformed roots from *Catharanthus roseus* and its relationship to tryptamine, ajmalicine and catharanthine accumulation during the culture cycle. *In Vitro Cell Dev Biol* 1994; 30P: 81-83.
186. Pennings, E. J. M., R. Verpoorte, O. J. M. Goddijn and J. H. C. Hoge. Purification of tryptophan decarboxylase from a *Catharanthus roseus* cell suspension culture. *J Chromatogr* 1989; 483(1): 311-318.
187. Schripsema, J., T. A. Van Beek, R. Verpoorte, C. Erkelens, P. Perera and C. Tibell. A reinvestigation of the stereochemistry of tubotaiwine using NMR spectroscopy. *J Nat Prod* 1987; 50(1): 89-101.
188. Kanamori-Fukuda, I., H. Ashihara and A. Komamine. Pyrimidine nucleotide biosynthesis in *Vinca rosea* cells: Changes in the activity of the de novo and salvage pathways during growth in a suspension culture. *J Exp Bot* 1981; 32(126): 69-78.
189. Tabgkongchitr, U. Extraction of organic compounds from leaves of *Vinca rosea* Linn. Thesis-MS-Chulalongkom Univ 1973; 1973: 42pp.
190. Smith, S. L., W. E. Jones, D. E. Dorman, E. E. Logsdon and G. H. Svoboda. Alkaloids of *Vinca rosea* L. (*Catharanthus roseus* G. Don). XXXIII. Isolation and characterization of new dimeric alkaloids. (Abstract). *Lloydia* 1974; 37(4): 645D.
191. Jones, W. E. and G. J. Cullinan. Vincadioline. Patent-US-3,887,565 1975; 4pp.
192. Hutchinson, C. R. Biosynthetic studies of antitumor indole alkaloids. *Annu Proc Phytochem Soc Eur* 1980; 17: 143-158.
193. Padasani, K. M., S. Singh and J. P. S. Sarin. Chromatographic estimation of vincalukoblastine in *Catharanthus roseus* Linn. *Indian J Pharm Sci* 1979; 41: 207-213.
194. Ramiah, N. Occurrence of liquid alkaloids in *Vinca rosea*. *J Indian Chem Soc* 1964; 41: 552.
195. Svoboda, G. H., M. Gorman and R. H. Tust. Alkaloids of *Vinca rosea* (*Catharanthus roseus*). XXV. Lochrovine, perimivine, vincoline, lochrovicine, lochrovicine and vincolidine. *Lloydia* 1964; 27: 203.

196. Aynilian, G. H., S. G. Weiss, G. A. Cordell, D. J. Abraham, F. A. Crane and N. R. Farnsworth. Catharanthus alkaloids. XXIX. Isolation and structure elucidation of vincoline. *J Pharm Sci* 1974; 63(4): 536-538.
197. Cuellar, A. Catharanthus roseus alkaloids. The structure of vincubine. *Rev Cubana Farm* 1976; 10: 19.
198. Song, K. M., S. W. Park, W. H. Hong, H. Lee, S. S. Kwak and J. R. Liu. Isolation of vindoline from *Catharanthus roseus* by supercritical fluid extraction. *Biotechnol Prog* 1992; 8(6): 583-586.
199. Kohl, W., H. Vogelmann and G. Hofle. Alkaloids from *Catharanthus roseus* tissue cultures. *Planta Med* 1980;39:283-284.
200. Moza, B. K. and J. Trojanek. New alkaloids from *Catharanthus roseus*. *Chem Ind (London)* 1962; 1962: 1425.
201. Davey, J. E., J. Van Staden and G. T. N. De Leeuw. Endogenous cytokinin levels and development of flower virescence in *Catharanthus roseus* infected with mycoplasmas. *Physiol Plant Pathol* 1981; 19: 193-200.

CONFLICT OF INTEREST REPORTED: NIL; SOURCE OF FUNDING: NONE REPORTED